

RATIONAL DECOMPOSITIONS OF COMPLEX MEROMORPHIC FUNCTIONS

ALAIN ESCASSUT AND EBERHARD MAYERHOFER

ABSTRACT. Let h be a complex meromorphic function. The problem of decomposing h in two different ways $P(f)$ and $Q(g)$ with f, g two other meromorphic functions and P, Q polynomials, was studied by C.-C. Yang, P. Li and K. H. Ha. Here we consider the problem when we replace the polynomials P, Q by rational functions F, G . Let $\deg(F)$ be the maximum degree of numerator and denominator of F . Assume some zeros $c_1, \dots, c_k$ of F' satisfy a pack of 5 conditions involving particularly $G(d) \neq F(c_j)$ and $D(d) \neq 0$ for every zero d of $C' - F(c_j)D'$, with $G = \frac{C}{D}$, ($j = 1, \dots, k$). First, we show that if f, g are entire functions such that $F(f) = G(g)$ then $k \deg(G) \leq \deg(F)$. Now, let $\gamma(D)$ be the number of distinct zeros of the denominator of G and assume that meromorphic functions f, g satisfying $F(f) = G(g)$, then $k \deg(G) \leq \deg(F) + k\gamma(D)$. When zeros $c_1, \dots, c_k$ of F' satisfy a stronger condition, then we show that $k \deg(G) \leq \deg(F) + k \min(\gamma(C), \gamma(D))$.

1. INTRODUCTION AND RESULTS

Let $\mathcal{A}(\mathbb{C})$ be the algebra of entire functions in $\mathbb{C}$ and $\mathcal{M}(\mathbb{C})$ be the field of meromorphic functions in $\mathbb{C}$. The decomposition of a function $h \in \mathcal{M}(\mathbb{C})$ was studied in [2] and [5] in order to find many cases when h cannot admit two different decompositions of the form $P(f)$ and $Q(g)$, with $P, Q \in \mathbb{C}[x]$ and $f, g \in \mathcal{M}(\mathbb{C})$. In [1], a similar study was made in a p-adic field K of characteristic $p \geq 0$, concerning meromorphic functions in all K , or meromorphic functions inside an "open" disk. In [7] the problem was considered in a p-adic field again, when P, Q lie not in $K[x]$ but in $K(x)$. Here we mean to consider this last problem in $\mathbb{C}$: given $h \in \mathcal{M}(\mathbb{C})$, can we find conditions excluding that h admits two different decompositions of the form $F(f)$ and $G(g)$, with $F, G \in \mathbb{C}(x)$? The Nevanlinna Second Main Theorem provides us with the main tool to deal with the problem, in this case as in the previous ones. We shall also use some basic algebraic consideration already made in [6].

Notation. Let $F = \frac{A}{B} \in \mathbb{C}(x)$ with $A, B \in \mathbb{C}[x]$ and $\gcd(A, B) = 1$. We put $\deg(F) = \max(\deg(A), \deg(B))$. Let $P \in \mathbb{C}[x]$. We denote by $\gamma(P)$ the number of distinct zeros of P . Given $F(x) = \frac{A}{B}$ with $A, B \in \mathbb{C}[x]$ and $\gcd(A, B) = 1$, we put $\lambda(F) = \min(\gamma(A), \gamma(B))$.

Definition. Let $\mathbb{E}$ be an algebraically closed field, let $F = \frac{A}{B}, G = \frac{C}{D} \in \mathbb{E}(x)$ with $A, B, C, D \in \mathbb{E}[x]$, $\gcd(A, B) = \gcd(C, D) = 1$. Let $c_1, \dots, c_k \in \mathbb{E}$ be zeros of F' . Then (F, G) will be said to *satisfy Condition M* for $c_1, \dots, c_k$ if:

- 1) C and D are monic,

2000 *Mathematics Subject Classification.* 30D35, 30D30, 39B32.

Key words and phrases. Nevanlinna Theory, Meromorphic Functions, Functional Equations.

- 2) $F(c_j) \neq F(c_l) \forall j \neq l$,
 - 3) if $\deg(C) = \deg(D)$ we assume $F(c_j) \neq 1 \forall j = 1, \dots, k$,
 - 4) for every $j = 1, \dots, k$ we have $G(d) \neq F(c_j)$ and $D(d) \neq 0$ for every zero d of $C' - F(c_j)D'$.
- Further, (F, G) will be said to *satisfy Condition M'* for $c_1, \dots, c_k$ if Conditions 1), 2), 3) are satisfied and if Condition 4) is replaced by Condition 4'):
- 4') for every $j = 1, \dots, k$ we have $G(d) \neq F(c_j)$ and $C(d)D(d) \neq 0$ for every zero d of $C' - F(c_j)D'$,

Theorem 1. *Let $F, G \in \mathbb{C}(x)$ satisfy Condition M for $c_1, \dots, c_k$ and let $p = \deg(F)$, $q = \deg(G)$. Assume that there exist two functions $f, g \in \mathcal{A}(\mathbb{C})$ such that $F(f) = G(g)$. Then $kq \leq p$.*

Theorem 2. *Let $F = \frac{A}{B}$, $G = \frac{C}{D} \in \mathbb{C}(x)$ with $\gcd(A, B) = \gcd(C, D) = 1$ and C, D monic, satisfy Condition M for $c_1, \dots, c_k$. Let $p = \deg(F)$, $q = \deg(G)$. Assume that there exist two functions $f, g \in \mathcal{M}(\mathbb{C})$ such that $F(f) = G(g)$. Then $kq \leq p(1 + (k\gamma(D)))$.*

Theorem 3. *Let $F = \frac{A}{B}$, $G = \frac{C}{D} \in \mathbb{C}(x)$ with $\gcd(A, B) = \gcd(C, D) = 1$ and C, D monic, satisfy Condition M' for $c_1, \dots, c_k$. Let $p = \deg(F)$, $q = \deg(G)$. Assume that there exist two functions $f, g \in \mathcal{M}(\mathbb{C})$ such that $F(f) = G(g)$. Then $kq \leq p(1 + k\lambda(G))$.*

Remarks. 1) Theorems 2 and 3 are trivial when C and D have same degree and have no multiple zero, and are more interesting when $\min(\deg(C), \deg(D))$ is much smaller than $\max(\deg(C), \deg(D))$.

2) Let $f(x) = \sin x$, $g(x) = \cos x$, $F(x) = \frac{x^2-1}{x^2}$, $G(x) = \frac{x^2}{x^2-1}$. Then $F(f) = G(g) = -\tan^2$, while $f, g \in \mathcal{A}(\mathbb{C})$. According to the notation of Theorem 1, we have $p = q = 2$, hence k must be at most 1. Actually this is right because $F'(x) = \frac{2}{x^3}$ has no zero.

Acknowledgement. The authors are very grateful to C.-C. Yang for letting us know the problem of the decomposition and the first results obtained in this domain by himself and his colleagues P. Li and H. K. Ha.

2. THE PROOFS

In order to prove the theorems we first need some notation and lemmas

Notation. Let ϕ, ψ be functions from $\mathbb{R}_+$ to $\mathbb{R}_+$. We put $\phi \lesssim \psi$ if there exists another function θ from $\mathbb{R}_+$ to $\mathbb{R}_+$ and a subset H of $\mathbb{R}_+$ of zero Lebesgue measure such that $\lim_{r \rightarrow \infty, r \notin H} \frac{\theta(r)}{\psi(r)} = 0$ and such that $\phi(r) \leq \psi(r) + \theta(r)$ (we notice that $\lesssim$ is a preorder).

In the same way, we put $\phi \sim \psi$ if there exists another function θ from $\mathbb{R}_+$ to $\mathbb{R}_+$ and a subset H of $\mathbb{R}_+$ of zero Lebesgue measure such that $\lim_{r \rightarrow \infty, r \notin H} \frac{\theta(r)}{\psi(r)} = 0$ and such that $\phi(r) = \psi(r) + \theta(r)$ (we notice that $\sim$ is an equivalence relation).

Lemma 1 is shown in [7] in a p-adic field of any characteristic $p \geq 0$. Actually it holds in any algebraically closed field:

Lemma 1. *Let $\mathbb{E}$ be an algebraically closed field, let $F = \frac{A}{B}$, $G = \frac{C}{D} \in \mathbb{E}(x)$ with*

$\gcd(A, B) = \gcd(C, D) = 1$, satisfy Condition M for $c_1, \dots, c_k$ and let $\deg(G) = q$. Then for each $j = 1, \dots, k$, we have factorizations of the form $F(x) - F(c_j) = (x - c_j)^{s_j} R_j(x)$ with $s_j \geq 2$, $R_j(c_j) \neq 0$ and $(G(x) - F(c_j))D(x) = \prod_{l=1}^q (x - b_{j,l})$ where the $b_{j,l}$ are qk distinct elements.

Lemma 2 is classical ([4],[6],[8]):

Lemma 2. Let $R \in \mathbb{C}(x)$ and let $f \in \mathcal{M}(\mathbb{C})$. Then $T(r, R(f)) \sim \deg(R)T(r, f)$.

Lemma 3 is basic:

Lemma 3. Let $\mathbb{E}$ be a field, let $F = \frac{A}{B} \in \mathbb{E}(x)$, let c be a zero of F' such that $F(c) \neq 0$. Then c is a zero of $(\frac{1}{F})'$, with the same multiplicity.

Lemma 4. Let $\mathbb{E}$ be an algebraically closed field, let $F = \frac{A}{B}$, $G = \frac{C}{D} \in \mathbb{E}(x)$ with $\gcd(A, B) = \gcd(C, D) = 1$, satisfy Condition M' for c_j , $1 \leq j \leq k$. Then $\frac{1}{F}, \frac{1}{G}$ also satisfy Condition M' for $c_1, \dots, c_k$.

Proof. By Lemma 3, $c_1, \dots, c_k$ are zeros of $(\frac{1}{F})'$. Next, Conditions 1), 2), 3) are obviously satisfied by $\frac{1}{F}, \frac{1}{G}$. So, we only have to show Condition 4'). Let u be a zero of $D' - \frac{1}{F(c_j)}C'$. First, u is a zero of $C' - F(c_j)D'$, hence by Condition 4') satisfied by F, G , we have $F(c_j) \neq G(u)$, therefore $\frac{1}{F(c_j)} \neq \frac{1}{G(u)}$. Moreover, by Condition 4') we have $D(u)C(u) \neq 0$ which is symmetric with regard to C and D . This ends the proof of Lemma 4. $\square$

Let us recall an opportune form of the Nevanlinna Second Main Theorem ([3], [4], [6], [8]):

Theorem N. Let $f \in \mathcal{M}(\mathbb{C})$. Let $b_1, \dots, b_n \in \mathbb{C}$. Then

$$(n-1)T(r, f) \lesssim \sum_{j=1}^n \overline{Z}(r, f - b_j) + \overline{N}(r, f).$$

Proof of the Theorems 1, 2 and 3. In the three Theorems, we put $F = \frac{A}{B}$, $G = \frac{C}{D}$ with $\gcd(A, B) = \gcd(C, D) = 1$ and C, D monic. By Lemma 1, for each $j = 1, \dots, k$ we can factorize $F(x) - F(c_j)$ in the form

$$(2.1) \quad (x - c_j)^{s_j} R_j(x)$$

with $s_j \geq 2$ and $R_j(c_j) \neq 0$ and similarly

$$(2.2) \quad (G(x) - F(c_j))D(x) = \prod_{l=1}^q (x - b_{j,l})$$

whereas the $b_{j,l}$ are qk distinct elements. Thus we obtain

$$(2.3) \quad F(f) - F(c_j) = (f - c_j)^{s_j} R_j(f) = \frac{1}{D(g)} \prod_{l=1}^q (g - b_{j,l}) = G(g) - F(c_j)$$

Now, by Theorem N we have

$$(2.4) \quad (qk-1)T(r, g) \lesssim \sum_{j=1}^k \sum_{l=1}^q \overline{Z}(r, g - b_{j,l}) + \overline{N}(r, g)$$

Thanks to (2), for each fixed j we obtain

$$(2.5) \quad \overline{Z}(r, (G(g) - F(c_j))D(g)) = \sum_{l=1}^q \overline{Z}(r, g - b_{j,l})$$

Then, by inserting g in (4) and in (5) we obtain

$$(2.6) \quad (qk - 1)T(r, g) \lesssim \sum_{j=1}^k \overline{Z}(r, (G(g) - F(c_j))D(g)) + \overline{N}(r, g)$$

Now, by Lemma 2 we have $T(r, g) \sim \frac{p}{q}T(r, f)$. Let $R_j(x) = \frac{A_j(x)}{B_j(x)}$: of course we have $\deg(A_j) \leq p - s_j$. By applying Lemma 2, we obtain

$$(2.7) \quad \overline{Z}(r, R_j(f)) + \overline{N}(r, f) = \overline{Z}(r, A_j(f)) + \overline{N}(r, f) \lesssim (p - s_j)T(r, f) + \overline{N}(r, f)$$

Now, assume the hypothesis of Theorem 1. Since f and g are entire and since on one hand, A and B have no common zero and on the other hand, C and D have no common zero, we can see that by (3), for each $j = 1, \dots, k$, any zero of $D(g)$ is a pole of $F(f)$ of same order, and therefore is a pole of $R_j(f)$ of same order. Therefore $\overline{Z}(r, R_j(f)D(g)) = \overline{Z}(r, R_j(f))$. Consequently, by (6) and (7), taking into account that f, g are entire, we can get

$$\begin{aligned} \overline{Z}(r, (G(g) - F(c_j))D(g)) &= \overline{Z}(r, (f - c_j)^{s_j} R_j(f)D(g)) \leq \\ &\leq \overline{Z}(r, (f - c_j)) + \overline{Z}(r, R_j(f)) \lesssim T(r, f) + (p - s_j)T(r, f) \end{aligned}$$

Summing up on j , by (6) we obtain $(qk - 1)T(r, g) \leq T(r, f)(k(1 + p) - \sum_{j=1}^k s_j)$. On the other hand, by Lemma 2 we have $T(r, f) \sim \frac{q}{p}T(r, g)$, therefore

$$(qk - 1)pT(r, f) \lesssim qT(r, f)(kp - \sum_{j=1}^k (s_j - 1))$$

Since $T(r, f)$ is unbounded, we can derive that $(qk - 1)p \leq q(kp - \sum_{j=1}^k (s_j - 1))$ hence $\sum_{j=1}^k (s_j - 1) \leq p$. But by hypothesis, $k \leq \sum_{j=1}^k (s_j - 1)$, thereby $kq \leq p$.

Next, assume the hypothesis of Theorem 2. We can find a constant $h \in \mathbb{C}$ such that $F(c_j) + h \neq 0 \forall j = 1, \dots, k$. Let $\Phi(x) = F(x) + h$, $\Psi(x) = G(x) + h$. Thus we have $\Phi(f) = \Psi(g)$, and Φ, Ψ satisfy the hypothesis of Theorem 2, in place of F, G , respectively. Consequently, we can assume that $F(c_j) \neq 0 \forall j = 1, \dots, k$, without loss of generality. Now, we have

$$\overline{Z}(r, R_j(f)D(g)) \leq \overline{Z}(r, R_j(f)) + \overline{Z}(r, D(g))$$

Consequently, by (6) and (7), we can get

$$(2.8) \quad \begin{aligned} \overline{Z}(r, (G(g) - F(c_j))D(g)) &= \overline{Z}(r, (f - c_j)^{s_j} R_j(f)D(g)) \leq \\ &\leq \overline{Z}(r, (f - c_j)) + \overline{Z}(r, R_j(f)) + \overline{Z}(r, D(g)) \end{aligned}$$

Let $t = \gamma(D)$ and let δ_u ($1 \leq u \leq t$) be the distinct zeros of D . We notice that $\overline{Z}(r, D(g)) \leq \sum_{u=1}^t \overline{Z}(r, g - \delta_u) \leq tT(r, g)$. Consequently, by (8) we obtain

$$\overline{Z}(r, (G(g) - F(c_j))D(g)) \lesssim T(r, f) + (p - s_j)T(r, f) + \gamma(D)T(r, g) \lesssim (p - (s_j - 1) + \frac{p\gamma(D)}{q})T(r, f)$$

Summing up on j , by (6) we obtain

$$(qk - 1)T(r, g) \leq T(r, f)(k(1 + p) - \sum_{j=1}^k s_j)$$

therefore

$$(qk - 1)pT(r, f) \lesssim_q T(r, f)(kp - \sum_{j=1}^k (s_j - 1) + \frac{kp\gamma(D)}{q})$$

Since $T(r, f)$ is unbounded, we can derive that $(qk - 1)p \leq q(kp - \sum_{j=1}^k (s_j - 1)) + kp\gamma(D)$ therefore $kq \leq q \sum_{j=1}^k (s_j - 1) \leq p(k\gamma(D) + 1)$.

Now, assume the hypothesis of Theorem 3. Thanks to Condition M', by Lemma 4 we can replace F , G by $\frac{1}{F}$ and $\frac{1}{G}$ respectively, because they also satisfy Condition M', hence we obtain $kq \leq p(k\gamma(C) + 1)$, consequently $kq \leq p(k\lambda(G) + 1)$. $\square$

REFERENCES

- [1] A. ESCASSUT AND C.-C. YANG, *The functional equation $P(f) = Q(g)$ in a p -adic field*, J. Number Theory, 105 (2004), pp. 344–360.
- [2] H. K. HA AND C.-C. YANG, *On the functional equation $P(f) = Q(g)$* , Advances in Complex Analysis and Applications, Kluwer Academic Pub., 2004.
- [3] W. K. HAYMAN, *Meromorphic functions*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1964.
- [4] I. LAINE, *Nevanlinna Theory and Complex Differential Equations*, de Gruyter Studies in Mathematics 15, de Gruyter, 1992.
- [5] P. LI AND C.-C. YANG, *Some further results on the unique range sets of meromorphic functions*, Kodai Math. J., 18 (1995), pp. 437–450.
- [6] A. I. MARKUSHEVICH, *Theory of functions of a complex variable. Vol. II*, Revised English edition translated and edited by Richard A. Silverman, Prentice-Hall Inc., Englewood Cliffs, N.J., 1965.
- [7] E. MAYERHOFER, *Rational decompositions of p -adic meromorphic functions*, Sci. Math. Jpn., 61 (2005), pp. 1–13.
- [8] R. NEVANLINNA, *Le théorème de Picard-Borel et la theorie des fonctions méromorphes*, Gauthiers-Villars, Paris, 1929.

UNIVERSITY OF VIENNA, DEPARTMENT OF MATHEMATICS, NORDBERGSTRASSE 15, 1090 VIENNA, AUSTRIA

E-mail address: eberhard.mayerhofer@univie.ac.at